


# Secrets of the Higgs Boson





M. E. Peskin  
HPNP 2015 Toyama  
February 2015

Only four years ago, in the fall of 2011, it was a popular theme for discussion among particle physicists that the Higgs boson did not exist.

Searches at the LHC had eliminated most of allowed range for the Higgs boson mass. Only a small corner remained in which the Higgs could hide.

Today, the situation could not be more different.





# Tests of qualitative properties predicted for the Higgs boson:

$\gamma\gamma$  decay mode ✓

$ZZ$  decay mode ✓

$WW$  decay mode ✓

$\tau^+\tau^-$  decay mode ✓


$b\bar{b}$  decay mode preliminary


$t\bar{t}$  coupling indirectly, through  $gg$

spin-parity  $0^+$  ✓


variables for the CMS  
spin analysis in

$$h \rightarrow ZZ^* \rightarrow \ell^+ \ell^- \ell^+ \ell^-$$





preference for Z to be longitudinally polarized




preference for Z decay planes to be parallel

The quantitative measure for used today for Higgs coupling values is the “signal strength”

$$\mu(A, B) = \frac{\sigma(A\bar{A} \rightarrow h)BR(h \rightarrow B\bar{B})}{(\text{SM expectation})}$$





PDG summary 2014

Today, received opinion in high-energy physics goes to the other extreme. The Higgs boson is discovered, it is Standard-Model-like, and its story is closed.

My view of the current situation is rather different.

I see the status of the Higgs boson as similar to that of the Cosmic Microwave Background in the late 1980's.

The Higgs boson is required by the Standard Model, and it is important to have its existence confirmed. Viewed at low precision, the Higgs boson must meet the simplest expectations. Only when we reach a required higher level of precision does the picture of the Higgs boson become richer and more instructive.

It is the purpose of this series of conferences to describe the patterns present in the Higgs coupling strengths that we will eventually search for in the experimental data.

In this lecture, I will give a broad-brush overview of this subject.

Today, we do not know whether electroweak spontaneous symmetry breaking is due to a single scalar field or to a more complicated, even strongly interacting, sector.


We do understand that this is a central -- in my opinion, **the** central -- question today in particle physics.

In a lecture at the 1981 Lepton-Photon Conference, Lev Okun described this structure with the symbol:

He emphasized that the search for and study of the Higgs boson was “problem #1” in particle physics.

Half of the Standard Model is governed by the gauge principle, which gives us a tight structure and perfect knowledge.

About the other hand, we are ignorant.




Gauge principle

/

Vacuum structure

Maxwell's  
equations  
Quantum  
numbers  
Parity  
violation  
W, Z boson  
Asymptotic  
freedom



W, Z mass  
Quark masses  
and mixings  
CP violation  
  
L, B violation  
  
Dark energy

Standard Model is  
all-powerful

Dark matter ?

Standard Model is  
impotent

It would seem that it should not be difficult to determine whether the Higgs sector contains one field or many, elementary or composite.

However, there is a barrier:

the “**Decoupling Theorem**” of Howard Haber

If the Higgs sector contains one light boson of mass

$$m_h = 125 \text{ GeV}$$

and many heavy particles with minimum mass  $M$ ,

the light boson has properties that **agree with the SM** predictions up to corrections of order

$$m_h^2 / M^2$$

Proof:

Integrate out the heavy fields. The result is the SM, plus a set of operators of minimum dimension 6.

Implication:

In most models of an extended Higgs sector or other new particles, the corrections to the Higgs couplings are at the **few-%** level. Precision measurement is needed to see these corrections.

However:

The **pattern of corrections** is different in different schemes for new physics models. **There is much to learn if we can see this pattern.**



Given the mass of the Higgs boson, the Standard Model makes a precise set of predictions for the couplings. These should be considered as **reference values** for precision measurements.

For a Higgs boson of mass 125 GeV, the prediction for the total width is  $\Gamma_h = 4.1 \text{ MeV}$

The branching fractions are predicted to be

$b\bar{b}$	58%	$\tau^+\tau^-$	6.3%	$\gamma\gamma$	0.23%
$WW^*$	21%	$c\bar{c}$	2.9%	$\gamma Z$	0.15%
$gg$	8.6%	$ZZ^*$	2.6%	$\mu^+\mu^-$	0.02%

Many decay modes of the Higgs will eventually be visible, and measurable. **F. Gianotti: “Thank you, Nature.”**

The study of the deviations from these predictions is guided by the idea that each Higgs coupling has **its own personality** and is guided by different types of new physics. This is something of a caricature, but, still, a useful one.

**fermion couplings** - multiple Higgs doublets

**gauge boson couplings** - Higgs singlets, composite Higgs

**$\gamma\gamma$ ,  $gg$  couplings** - heavy vectorlike particles

**$tt$  coupling** - top compositeness

**$hhh$  coupling** (large deviations) - baryogenesis

In a model with two Higgs doublets, the physical states are mixtures of the two fields

$$\text{mixing angle} \quad \begin{array}{ll} \alpha : & h^0, H^0 \\ \beta : & \pi^0, A^0 \end{array} \quad \begin{array}{l} \pi^\pm, H^\pm \end{array} \quad \tan \beta = v_u/v_d$$

Then the coupling modifications are


$$g(b\bar{b}) = -\frac{\sin \alpha}{\cos \beta} \frac{m_b}{v} \quad g(c\bar{c}) = \frac{\cos \alpha}{\sin \beta} \frac{m_c}{v}$$

In full models such as SUSY, the two angles are not independent. In fact, typically,

$$-\frac{\sin \alpha}{\cos \beta} = 1 + \mathcal{O}\left(\frac{m_Z^2}{m_A^2}\right)$$

so that


$$\frac{g_{hbb}}{g_{\text{SM}bb}} = \frac{g_{h\tau\tau}}{g_{\text{SM}\tau\tau}} \simeq 1 + 40\% \left( \frac{200 \text{ GeV}}{m_A} \right)^2$$




Kanemura, Tsumura, Yagyu, Yokoya

Loops with b,t squarks and gluinos can also modify this vertex, especially at large  $\tan \beta$ .

Haber, Herrero,  
Logan, Penaranda,  
Rigolin, Temes




$$\Gamma(h \rightarrow b\bar{b})$$



Cahill-Rowley, Hewett, Ismail, Rizzo

$$\Gamma(h \rightarrow \tau^+ \tau^-)$$



Cahill-Rowley, Hewett, Ismail, Rizzo

The coupling of the Higgs boson to vector bosons is similarly simple in the SM:

$$g(hVV) = \frac{2m_V^2}{v}$$

Corrections from models with an extended Higgs sector are usually small, since it is the lightest Higgs that has the largest vacuum expectation value. In SUSY,

$$g(hVV) = 1 + \mathcal{O}\left(\frac{m_Z^4}{m_A^4}\right)$$



Still, the  $hWW$  and  $hZZ$  coupling can obtain corrections from a number of sources outside the SM.

Mixing of the Higgs with a singlet gives corrections

$$g(hVV) \sim \cos \phi \sim (1 - \phi^2/2)$$

These might be most visible in the  $hVV$  couplings. Similarly, field strength renormalization of the Higgs can give 1% level corrections (Craig and McCullough).


If the Higgs is a composite Goldstone boson, these couplings are corrected by ( $f \sim 1$  TeV)

$$g(hVV) = (1 - v^2/f^2)^{1/2} \approx 1 - v^2/2f^2 \approx 1 - 3\%$$

The decays

$$h \rightarrow gg, \quad h \rightarrow \gamma\gamma, \quad h \rightarrow \gamma Z^0$$

proceed through loop diagrams.



The loops are dominated by **heavy particles** that the Higgs boson cannot decay to directly.

However, again, **decoupling** puts a restriction:

Only the heavy particles of the SM, that is,  **$t$ ,  $W$ ,  $Z$** , get 100% of their mass from the Higgs. For BSM particles such as  $\tilde{t}$  or  $T$ , the contribution to these loops is proportional to the fraction of their mass that comes from the Higgs vev.

Then, for example, a vectorlike T quark contributes

$$g(hgg)/SM = 1 + 2.9\% \left( \frac{1 \text{ TeV}}{m_T} \right)^2$$


$$g(h\gamma\gamma)/SM = 1 - 0.8\% \left( \frac{1 \text{ TeV}}{m_T} \right)^2$$

A complete model will have several new heavy states, and mixing of these with the SM top quark. For example, for the “Littlest Higgs” model

$$g(hgg)/SM = 1 - (5 - 9\%)$$


$$g(h\gamma\gamma)/SM = 1 - (5 - 6\%)$$

$\tan\beta=60$



Carena, Gori, Shah, Wagner

# Littlest Higgs model




Han, Logan, McElrath, Wang


In composite Higgs models, the shifts in the  $\gamma\gamma$  and  $gg$  partial widths come both from the modification of the top quark coupling and from the contributions of heavy vectorlike particles.

These effects are disentangled by direct measurement of the Higgs coupling to  $t\bar{t}$ .

Substantial effects are expected in models of partial top compositeness, as in Randall-Sundrum models.



Malm, Neubert, Schmell



Kanemura, Kaneta, Machida, Shindou




The Higgs self-coupling is a special case in this story.

Whereas we can expect the other Higgs couplings to be measured at the percent level, the  $hhh$  coupling is much more difficult to access.

The current expectation for ILC is a 16% measurement with 2000/fb at 1 TeV. Though it is expected that the LHC will have some sensitivity to this coupling, this has not yet been demonstrated (see ATL-PHYS-PUB-2014-019).


However, order-1 deviations in the  $hhh$  coupling are expected in some scenarios, in particular, in models of baryogenesis at the electroweak scale. These may be the only models of baryogenesis testable with accelerator data.

contours of fixed hhh coupling



Grojean, Servant, Wells

$\lambda_3$  vs  $m_h$  for  $\xi > 1$




Noble and Perelstein


Putting all of these effects together, we find patterns of deviations from the SM predictions that are different for different schemes of new physics.

For example:

## SUSY



## Composite Higgs



Kanemura, Tsumura, Yagyu, Yokoya

There is one more question on the theory side that we must address:

In order to measure a deviation from the Standard Model expectation, we must be able to compute the Standard Model reference values to better than 1% accuracy. Will this actually be possible ?

This has been questioned in the literature, especially for  $\Gamma(h \rightarrow b\bar{b})$ , for which a very accurate value of the b quark mass is needed.

There are two types of contributions to the theoretical error on SM predictions:

error from uncalculated orders of perturbation theory

error from uncertainty in input parameters (  $m_b, \alpha_s$  )

For the errors of the first type, the situation is quite good. These theoretical uncertainties are currently

0.3% for Higgs couplings to quarks

3 % for Higgs couplings to  $gg$

1 % for Higgs couplings to  $WW, ZZ$

1 % for Higgs coupling to  $\gamma\gamma$

Among the most impressive theoretical efforts are

Baikov, Chetyrkin, Kuhn:  $g(hb\bar{b})$  to  $\mathcal{O}(\alpha_s^4)$

Baikov, Chetyrkin,

Schreck and Steinhauser:  $g(hgg)$  to  $\mathcal{O}(\alpha_s^4)$

Actis, Passarino, Sturm, Uccirati:  $g(hgg)$  to  $\mathcal{O}(\alpha\alpha_s)$

Improvement of the current results to 0.1% accuracy is possible, but it will require dedicated effort.

For the dependence on input parameters, the situation is easier to quantify. The most important dependences of Higgs coupling predictions are  
(  $\delta_A = \Delta\Gamma(A)/\Gamma(A)$  ):

$$\delta_b = 1. \cdot \delta m_b(10) \oplus (-0.28) \cdot \delta\alpha_s(m_Z)$$

$$\delta_c = 1. \cdot \delta m_c(3) \oplus (-0.80) \cdot \delta\alpha_s(m_Z)$$

$$\delta_g = 1.2 \cdot \delta\alpha_s(m_Z)$$

We need to know the inputs in this table to the 0.1% level.



Many of the best determinations of these quantities now come from Lattice QCD. Mackenzie, LePage, and I projected the errors from Lattice QCD ten years into the future and estimated:

	$\delta m_b(10)$	$\delta \alpha_s(m_Z)$	$\delta m_c(3)$	$\delta_b$	$\delta_c$	$\delta_g$
current errors [10]	0.70	0.63	0.61	0.77	0.89	0.78
+ PT	0.69	0.40	0.34	0.74	0.57	0.49
+ LS	0.30	0.53	0.53	0.38	0.74	0.65
+ LS <sup>2</sup>	0.14	0.35	0.53	0.20	0.65	0.43
+ PT + LS	0.28	0.17	0.21	0.30	0.27	0.21
+ PT + LS <sup>2</sup>	0.12	0.14	0.20	0.13	0.24	0.17
+ PT + LS <sup>2</sup> + ST	0.09	0.08	0.20	0.10	0.22	0.09
ILC goal				0.30	0.70	0.60

relative errors in percent

The partial widths to WW, ZZ also depend strongly on the mass of the Higgs boson:

$$\delta_W = 6.9 \cdot \delta m_h , \quad \delta_Z = 7.7 \cdot \delta m_h$$

This is a 0.2% uncertainty for  $\Delta m_h = 30 \text{ MeV}$ .

This is the primary motivation (in my opinion) for a very accurate Higgs mass measurement.

So, if there is new physics beyond the Standard Model at the TeV scale, we expect to see deviations of the Higgs couplings from their Standard Model expectations at least at the few-percent level.

These deviations will form a pattern that will give us information on the nature of the new physics.

These ideas give strong motivation for a program of precision measurements of the Higgs boson couplings.

The goal should be to measure the individual partial widths to an accuracy of 1%, and better if possible.



This requires a comprehensive program on Higgs production and decay processes, such that the partial widths can be extracted by a combination of  $\mu(A, B)$  and cross section measurements.

It would be best if the experiments were also highly sensitive to invisible and exotic Higgs decays, which might contribute to  $\Gamma_h$  and also signal new physics in their own right.

We will learn much about the Higgs boson from its study at the LHC over the next 20 years.

However, the LHC cannot fulfill the goals of the program I have outlined.

The most important reasons for this are made clear by looking at the current evidence for the Higgs boson in its various decay channels.



**ATLAS  $H \rightarrow WW^*$**   
 $\sqrt{s} = 8 \text{ TeV}, 20.3 \text{ fb}^{-1}$

Obs  $\pm$  stat  
Exp  $\pm$  syst

Higgs  
WW  
Misid  
VV  
DY  
Top










# ATLAS EXPERIMENT

Run: 204153

Event: 35369265


2012-05-30 20:31:28 UTC











CMS, 4.9 fb<sup>-1</sup> at 7 TeV, 19.7 fb<sup>-1</sup> at 8 TeV







## CMS projections for the measurement accuracy of Higgs couplings at LHC and HL-LHC :

L (fb <sup>-1</sup> )	H → $\gamma\gamma$	H → WW	H → ZZ	H → bb	H → $\tau\tau$	H → Z $\gamma$	H → inv.
300	[6, 12]	[6, 11]	[7, 11]	[11, 14]	[8, 14]	[62, 62]	[17, 28]
3000	[4, 8]	[4, 7]	[4, 7]	[5, 7]	[5, 8]	[20, 24]	[6, 17]

CMS assumed that these accuracies are limited mainly by statistics and by theory uncertainties in the total cross sections. However, really, the limiting factor will be the **10:1 ratio of background to signal** in the best signal regions.

(For a comparison of CMS and ATLAS projections, see my paper [arXiv:1312.4974](https://arxiv.org/abs/1312.4974).)

One important special case should be noted:


The modes  $h \rightarrow \gamma\gamma$  and  $h \rightarrow 4\ell$  are visible in the total cross section, with very similar selections.

It should be possible to design an analysis in which measurement the ratio of the production rates, in the same specified region of  $\eta$ , is independent of the Higgs production cross section and is limited by statistics only.

ATLAS estimated the ultimate error on this ratio as 3.6%. This is probably an overestimate. The pure statistics limit (50% efficiency) is about 2% for 3000/fb.


There is a proposed accelerator capable of meeting these goals that is studied thoroughly, designed at the level its TDR, and ready for construction.

This is the International Linear Collider (ILC).




Here are a few snapshots from the physics expectations for the ILC.

$P(e^-, e^+) = (-0.8, 0.2)$ ,  $M_h = 125 \text{ GeV}$






$m_h$  to 30 MeV using a recoil technique




$$e^+e^- \rightarrow Z^0 \quad h^0$$


$$\rightarrow \mu^+\mu^- \quad b\bar{b}$$




**Data**




**$h \rightarrow \text{others}$**




**SM BG**




**$h \rightarrow bb$**




**$h \rightarrow cc$**




**$h \rightarrow gg$**








# Projected Higgs Coupling Precision, Model-Independent Fit




# Sensitivity to the extended Higgs sector of SUSY


LHC 300/fb




LHC 3000/fb



ILC TDR



ILC 20 yr



Cahill-Rowley, Hewett, Ismail, Rizzo

The precision study of the Higgs boson will be one of the next great adventures in particle physics.

The Higgs boson has many secrets that are still hidden. But it is within our power to find them out.