

Radiative corrections to electroweak parameters in the Higgs Triplet Model and implication with the recent Higgs boson searches at LHC

Kei Yagyu (Univ. of Toyama)

S. Kanemura, K. Yagyu, arXiv: 1201.6287 [hep-ph]

Toyama, Feb. 20th 2012

Introduction

- **The Higgs sector is unknown.**

- Minimal? or Non-minimal?
- The Higgs boson search is underway at the LHC.
The Higgs boson mass is constrained to be
 $115 \text{ GeV} < m_h < 127 \text{ GeV}$ or $m_h > 600 \text{ GeV}$.
- By the combination with electroweak precision data at the LEP,
we may expect that a light Higgs boson exists.

- **There are phenomena which cannot explain in the SM.**

- Tiny neutrino masses
- Existence of dark matter
- Baryon asymmetry of the Universe

- **New physics may explain these phenomena above the TeV scale.**

- Extended Higgs sectors are often introduced.


Physics of the Higgs sector  New physics beyond the SM

Explanation by extended Higgs sectors

- **Tiny neutrino masses**

- The type II seesaw model
- Radiative seesaw models

(e.g. Zee model)




- **Dark matter**

- Higgs sector with the discrete symmetry

- **Baryon asymmetry of the Universe**

- Electroweak baryogenesis



Introduce extended Higgs sectors

SU(2) doublet Higgs + Singlet [U(1)_{B-L} model],

SU(2) doublet Higgs + Doublet [Inert doublet model],

SU(2) doublet Higgs + Triplet [Type II seesaw model], etc...

How we can constrain these possibilities?

Constraint from the rho parameter

- ★ The experimental value of the rho parameter is quite close to unity.
- ★ Prediction of the rho parameter strongly depends on the structure of the extended Higgs sector.

$$\rho_{\text{exp}} \sim 1$$

$$\rho \equiv \frac{m_W^2}{m_Z^2 \cos^2 \theta_W} = \frac{\sum_i [T_i(T_i + 1) - Y_i^2] v_i^2}{\sum_i 2Y_i^2 v_i^2}$$

Y_i : hypercharge

T_i : isospin

v_i : VEV

Model with $\rho = 1$ at the tree level

- The Standard Model
- Models with Multi-doublet fields (with singlets)

The **custodial SU(2) symmetry** exists in the kinetic term.

Model with $\rho \neq 1$ at the tree level


- Model with a Y=0 Triplet field
- Model with a Y=1 Triplet field
- Models with larger isospin rep.

The **custodial SU(2) symmetry** does not exist in the kinetic term.

How the ρ parameter is calculated in both classes of models at the loop level.

Models with $\rho = 1$ at the tree level

The electroweak parameters are described by the 3 (+2) input parameters.



$$g, g', v + (Z_B, Z_W)$$

We can choose α_{em} , G_F and m_Z as the 3 input parameters.

$$\alpha_{\text{em}}(m_Z) = 128.903 \pm 0.0015$$

$$G_F = 1.16637 \pm 0.00001 \text{ GeV}^{-2}$$

$$m_Z = 91.1876 \pm 0.0021 \text{ GeV}$$

The other parameters can be written in terms of the above 3 inputs.

$$v^2 = \frac{1}{\sqrt{2}G_F}$$

$$s_W^2 = 1 - \frac{m_W^2}{m_Z^2}$$

$$m_W^2 s_W^2 = \frac{\pi \alpha_{\text{em}}}{\sqrt{2}G_F}$$


The deviation of the tree level relation can be expressed by Δr :

$$m_W^2 s_W^2 = \frac{\pi \alpha_{\text{em}}}{\sqrt{2}G_F} (1 + \Delta r)$$

Scalar boson and fermion loop contributions to Δr

$$\Delta r = -\frac{\delta G_F}{G_F} + \frac{\delta \alpha_{\text{em}}}{\alpha_{\text{em}}} - \frac{\delta m_W^2}{m_W^2} - \boxed{\frac{\delta s_W^2}{s_W^2}}$$

$$\Pi_T^{XY}(p^2) = \text{Diagram: } \text{wavy line } X \text{ with momentum } p \text{ enters a circle loop, and wavy line } Y \text{ exits.}$$




On-shell
renormalization
scheme

$$\Delta r_{\rho=1} = \frac{\Pi_T^{WW}(0) - \Pi_T^{WW}(m_W^2)}{m_W^2} + \frac{d}{dp^2} \Pi_T^{\gamma\gamma}(p^2) \Big|_{p^2=0} + \frac{2s_W}{c_W} \frac{\Pi_T^{\gamma Z}(0)}{m_Z^2} + \delta_{VB}$$

$$- \frac{c_W^2}{s_W^2} \left[\frac{\Pi_T^{ZZ}(m_Z^2)}{m_Z^2} - \frac{\Pi_T^{WW}(m_W^2)}{m_W^2} \right] \sim \rho - 1 = \alpha_{\text{em}} T$$

$$\simeq \frac{1}{16\pi^2} \left[\frac{(m_t - m_b)^2}{m_W^2} + \frac{(m_{H^+} - m_A)^2}{m_W^2} - \ln m_h^2 \right]$$

(In the case of the two Higgs doublet model)




Dependence of the **quadratic mass splitting** among particles in the same isospin multiplet appears in the T (rho) parameter.

Peskin, Wells (2001);

Grimus, Lavoura, OGREID, OSLAND (2008);

Kanemura, Okada, Taniguchi, Tsumura (2011).



Previous works and motivation of our work

- In models with $\rho \neq 1$ at the tree level, the renormalization scheme is different from that in models with $\rho=1$ at the tree level.
- In the model with the $Y=0$ Higgs triplet field, one-loop corrections to the electroweak parameters have been studied in Blank, Hollik (1997), Chen, Dawson (2004) etc.

NEW


- **We first study one-loop corrections to the electroweak precision parameters in the $Y = 1$ Higgs Triplet Model which is introduced in the type II seesaw mechanism. We then discuss how the model can be constrained by the data.**
- **Under this constraint, we discuss the implication with the recent Higgs boson searches at the LHC.**

The type II seesaw model (The Y=1 Higgs Triplet Model)


The Higgs triplet field Δ ($Y = 1$) is added to the SM.

$$\mathcal{L}_{\text{typeII}} = h_{ij} \overline{L}_L^{ci} \cdot \Delta L_L^j + \mu \Phi \cdot \Delta^\dagger \Phi + \dots$$

*Cheng, Li (1980);
Schechter, Valle, (1980);
Magg, Wetterich, (1980);
Lazarides, Shafi, Wetterich, (1981);
Mohapatra, Senjanovic, (1981).*



$$(m_\nu)_{ij} = h_{ij} \frac{\mu \langle \phi^0 \rangle^2}{M_\Delta^2} = h_{ij} v_\Delta$$




M_Δ : Mass of triplet scalar boson.
 v_Δ : VEV of the triplet Higgs

When we consider the TeV scale M_Δ , the $L\#$ violating coupling μ has to be of $O(10^{-10})$ GeV.

Models with $\rho \neq 1$ at the tree level

The electroweak parameters are described by the **4** (+2) input parameters.




$$g, g', v_\Phi, v_\Delta + (Z_B, Z_W)$$

Blank, Hollik (1997)

We can choose $\alpha_{\text{em}}, G_F, m_Z$ and $\hat{s}_W^2$ as the input parameters.

$\hat{s}_W^2$ is defined by the effective Zee vertex:

$$\mathcal{L} = \bar{e} \frac{g}{2\hat{c}_W} (v_e \gamma_\mu - a_e \gamma_\mu \gamma_5) e Z^\mu$$



$$1 - 4\hat{s}_W^2(m_Z) = \frac{\text{Re}(v_e)}{\text{Re}(a_e)}$$

$$\hat{s}_W^2 = 0.23146 \pm 0.00012$$

The other parameters are determined as:

$$v^2 = v_\Phi^2 + 2v_\Delta^2 = \frac{1}{\sqrt{2}G_F}$$

$$m_W^2 = \frac{\pi\alpha_{\text{em}}}{\sqrt{2}G_F\hat{s}_W^2}$$



$$v_\Delta^2 = \frac{\hat{s}_W^2(1 - \hat{s}_W^2)}{2\pi\alpha_{\text{em}}} m_Z^2 - \frac{\sqrt{2}}{4G_F}$$

Radiative corrections to EW parameters in models with $\rho \neq 1$ at the tree level

In the model with $\rho \neq 1$: s_W^2 is **an independent parameter**.

→ **Additional renormalization condition is necessary.**

Blank, Hollik (1997)



$$\frac{\delta \hat{s}_W^2}{\hat{s}_W^2} = \frac{\hat{c}_W}{\hat{s}_W} \frac{\Pi_T^{\gamma Z}(m_Z^2)}{m_Z^2} - \delta'_V$$

$$\Delta r_{\rho \neq 1} = \frac{\Pi_T^{WW}(0) - \Pi_T^{WW}(m_W^2)}{m_W^2} + \frac{d}{dp^2} \Pi_T^{\gamma\gamma}(p^2) \Big|_{p^2=0} + \frac{2\hat{s}_W}{\hat{c}_W} \frac{\Pi_T^{\gamma Z}(0)}{m_Z^2} + \delta_{VB} + \frac{\hat{c}_W}{\hat{s}_W} \frac{\Pi_T^{\gamma Z}(m_Z^2)}{m_Z^2} + \delta'_V$$

$$\simeq \frac{1}{16\pi^2} [\ln m_t^2 + \ln m_{\Delta\text{-like}}^2 + \ln m_h^2]$$

By the renormalization of δs_W^2 , quadratic dependence of the mass splitting disappear.

$$\rho = \frac{\pi\alpha_{\text{em}}}{\sqrt{2}G_F m_Z^2 \hat{s}_W^2 \hat{c}_W^2} (1 + \Delta r)$$

$$m_W^2 = \frac{\pi\alpha_{\text{em}}}{\sqrt{2}G_F \hat{s}_W^2} (1 + \Delta r)$$

Higgs potential in the HTM

Higgs potential

$$V = m^2 \Phi^\dagger \Phi + M^2 \text{Tr}(\Delta^\dagger \Delta) + [\mu \Phi^T i \tau_2 \Delta^\dagger \Phi + \text{h.c.}] \\ + \lambda_1 (\Phi^\dagger \Phi)^2 + \lambda_2 [\text{Tr}(\Delta^\dagger \Delta)]^2 + \lambda_3 \text{Tr}[(\Delta^\dagger \Delta)^2] \\ + \lambda_4 (\Phi^\dagger \Phi) \text{Tr}(\Delta^\dagger \Delta) + \lambda_5 \Phi^\dagger \Delta \Delta^\dagger \Phi.$$

$$\Phi = \begin{bmatrix} \varphi^+ \\ \frac{1}{\sqrt{2}}(\varphi + v + i\chi) \end{bmatrix} \\ \Delta = \begin{pmatrix} \frac{\Delta^+}{\sqrt{2}} & \Delta^{++} \\ \Delta^0 & -\frac{\Delta^+}{\sqrt{2}} \end{pmatrix} \\ \Delta^0 = \frac{1}{\sqrt{2}}(\delta + v_\Delta + i\eta)$$

Mass eigenstates: (SM-like) **h**, (Triplet-like) **H^{±±}, H[±], H, A**


Mass spectrum:

$$m_h^2 \simeq 2\lambda_1 v^2$$


$$M_\Delta^2 \equiv \frac{v^2 \mu}{\sqrt{2} v_\Delta}$$

$$m_{H^{++}}^2 \simeq M_\Delta^2 - \frac{v^2}{2} \lambda_5 \\ m_{H^+}^2 \simeq M_\Delta^2 - \frac{v^2}{4} \lambda_5 \\ m_A^2 \simeq m_H^2 = M_\Delta^2$$

Case I ($\lambda_5 > 0$)



Case II ($\lambda_5 < 0$)



We discuss the constraint from the electroweak precision data
In both **Case I** and **Case II**.


Prediction to the rho parameter at the 1-loop level

Kanemura, Yagyu, arXiv: 1201.6287 [hep-ph]

$$\Delta m = m_{H^{++}} - m_{H^+}$$

Case I: $m_{H^{++}} = 150$ GeV, $m_h = 125$ GeV, $\tan\alpha = 0$

Case II: $m_A = 150$ GeV, $m_h = 125$ GeV, $\tan\alpha = 0$



v_Δ is calculated according to the tree level relation:

$$v_\Delta^2 = \frac{\hat{s}_W^2(1 - \hat{s}_W^2)}{2\pi\alpha_{\text{em}}} m_Z^2 - \frac{\sqrt{2}}{4G_F}$$

In Case I with $m_{H^{++}} = 150$ GeV, $100 \text{ GeV} < |\Delta m| < 400 \text{ GeV}$ and $3 \text{ GeV} < v_\Delta < 8 \text{ GeV}$ is allowed. Case II is highly constrained by the rho parameter data.


Prediction to the W boson mass at the 1-loop level

$$m_{H^{++}} = 150 \text{ GeV}$$


$$m_{H^{++}} = 300 \text{ GeV}$$

$$\Delta m = m_{H^{++}} - m_{H^+}$$


Case I: $m_{H^{++}} = 150 \text{ GeV}$, $m_h = 125 \text{ GeV}$, $\tan\alpha = 0$




Case I: $m_{H^{++}} = 300 \text{ GeV}$, $m_h = 125 \text{ GeV}$, $\tan\alpha = 0$



Case I




Case II: $m_A = 150 \text{ GeV}$, $m_h = 125 \text{ GeV}$, $\tan\alpha = 0$




$$m_A = 150 \text{ GeV}$$

Case II: $m_A = 300 \text{ GeV}$, $m_h = 125 \text{ GeV}$, $\tan\alpha = 0$



$$m_A = 300 \text{ GeV}$$

Case II



In Case I, by the effect of the mass splitting, there are allowed regions .
Case II is highly constrained by the data.

Heavy mass limit

$$\xi = m_{H^{++}}^2 - m_{H^+}^2$$

Case I


Case II

$$\rho_{\text{SM}} \simeq 1 + \frac{\hat{c}_W^2}{\hat{c}_W^2 - \hat{s}_W^2} \frac{N_c \sqrt{2}}{16\pi^2} G_F m_t^2$$

$$\rho_{\text{HTM}} \simeq 1 + \frac{g^2 N_c}{16\pi^2} \ln m_t$$


$$\Delta\rho \equiv \rho - \rho_{\text{SM}}(m_h^{\text{ref}} = 125 \text{ GeV})$$

$$\Delta\rho^{\text{exp}} = 0.000632 \pm 0.000621$$




When we take heavy mass limit, loop effects of the triplet-like scalar bosons disappear. Even in such a case, the prediction does not coincide with the SM prediction.

Higgs $\rightarrow$ two photon decay




SM contribution

+




Triplet-like scalar loop contribution


Case I: $m_{H^{++}} = 150$ GeV, $m_h = 125$ GeV, $v = 6.7$ GeV, $\tan\alpha = 0$



Case I, $m_{H^{++}} = 300$ GeV, $m_h = 125$ GeV, $v_\Delta = 6.7$ GeV, $\tan\alpha = 0$



Case I




$$R_{\gamma\gamma} \equiv \frac{\Gamma(h \rightarrow \gamma\gamma)_{\text{HTM}}}{\Gamma(\phi_{\text{SM}} \rightarrow \gamma\gamma)_{\text{SM}}}$$

The decay rate of $h \rightarrow \gamma\gamma$ is around half in the HTM compared with that in the SM.

Summary

- Electroweak precision data (ρ , m_W , ...) can be constrained to the structure of extended Higgs sectors.
- In models with $\rho \neq 1$ at the tree level, **4 input parameters (α_{em} , G_F , m_Z and $\hat{s}_W^2$) instead of 3 ones (α_{em} , G_F and m_Z)** are necessary to describe the electroweak parameters. $\rightarrow$ An additional renormalization condition is required to renormalize the electroweak parameters.
- Case II is strongly constrained by the electroweak precision data, on the other hand in Case I with **$m_{H^{++}} \sim 150$ GeV, $|\Delta m| \sim \text{several } 100$ GeV and $v\Delta \sim O(1)$ GeV is favored by the data.**
- In the allowed parameter regions by the data, the decay rate of $h \rightarrow \gamma\gamma$ is around 50% in the HTM compared to that in the SM.

On-shell renormalization scheme



On-shell renormalization conditions

$$\left. \begin{array}{c} p \rightarrow \\ \text{wavy line} \\ W, Z \end{array} \right\} \text{IPI diagram} \left. \begin{array}{c} \text{wavy line} \\ W, Z \end{array} \right\} \left|_{p^2 = m_{W,Z}^2} = 0$$

$$\left. \begin{array}{c} p \rightarrow \\ \text{wavy line} \\ \gamma \end{array} \right\} \text{IPI diagram} \left. \begin{array}{c} \text{wavy line} \\ Z \end{array} \right\} \left|_{p^2 = 0} = 0$$

$$\frac{d}{dp^2} \left[\begin{array}{c} p \rightarrow \\ \text{wavy line} \\ \gamma \end{array} \right\} \text{IPI diagram} \left. \begin{array}{c} \text{wavy line} \\ \gamma \end{array} \right\} \right]_{p^2 = 0} = 0$$

$$\left. \begin{array}{c} e^- \text{ with } p_1 \\ e^+ \text{ with } p_2 \end{array} \right\} \text{IPI diagram} \left. \begin{array}{c} \text{wavy line} \\ \gamma \end{array} \right\} \left|_{\substack{p^2 = 0, \\ \not{p}_1 = \not{p}_2 = m_e}} = 0$$

From these 5 conditions, 5 counter terms (δg , $\delta g'$, δv , δZ_B , δZ_W) are determined.

Radiative corrections to the EW parameters

The deviation form $m_W^2 s_W^2 = \frac{\pi \alpha_{\text{em}}}{\sqrt{2} G_F}$
can be parametrized as:

$$m_W^2 s_W^2 = \frac{\pi \alpha_{\text{em}}}{\sqrt{2} G_F} (1 + \Delta r)$$

$$\Delta r = -\frac{\delta G_F}{G_F} + \frac{\delta \alpha_{\text{em}}}{\alpha_{\text{em}}} - \frac{\delta s_W^2}{s_W^2} - \frac{\delta m_W^2}{m_W^2}$$

From the renormalization conditions;

$$\frac{\delta \alpha_{\text{em}}}{\alpha_{\text{em}}} = \frac{d}{dp^2} \Pi_T^{\gamma\gamma}(p^2) \Big|_{p^2=0} + \frac{2s_W}{c_W} \frac{\Pi_T^{\gamma Z}(0)}{m_Z^2}$$

$$\frac{\delta G_F}{G_F} = -\frac{\Pi_T^{WW}(0)}{m_W^2} - \delta_{VB}$$

$$\frac{\delta m_W^2}{m_W^2} = \frac{\Pi_T^{WW}(m_W^2)}{m_W^2}$$

In models with $\rho = 1$ at the tree level, s_W^2 is the dependent parameter.
Therefore, the counter term for δs_W^2 is given by the other conditions.

$$s_W^2 = 1 - \frac{m_W^2}{m_Z^2} \quad \Rightarrow \quad \frac{\delta s_W^2}{s_W^2} = \frac{c_W^2}{s_W^2} \left[\frac{\Pi_T^{ZZ}(m_Z^2)}{m_Z^2} - \frac{\Pi_T^{WW}(m_W^2)}{m_W^2} \right]$$


This part represents the violation of the custodial symmetry
by the sector which is running in the loop.

Phenomenology of HTM with the mass splitting at the LHC




Aoki, Kanemura, Yagyu, Phys. Rev. D, in press (2011)

Case II

Cascade decays of the Δ -like scalar bosons become important.



Case I




Transvers mass

$$M_T^2 = (\cancel{E}_T + p_T)^2 \simeq 2|\cancel{E}_T||p_T|(1 - \cos \varphi)$$


By using the M_T distribution, we may reconstruct the mass spectrum of Δ -like scalar bosons.
 → We would test the Higgs potential in the HTM.

$m_h = 700$ GeV case


Case I: $m_{H^{\pm}} = 150$ GeV, $m_h = 700$ GeV, $\tan\alpha = 0$




Case I: $m_{H^{\pm}} = 300$ GeV, $m_h = 700$ GeV, $\tan\alpha = 0$




Case II: $m_A = 150$ GeV, $m_h = 700$ GeV, $\tan\alpha = 0$



Case II: $m_A = 300$ GeV, $m_h = 700$ GeV, $\tan\alpha = 0$




$$m_h = 125 \text{ GeV}, \tan\alpha = 2v_\Delta/v, v_\Delta = 0$$



Exotic Higgs Φ^{++} : EPS Results

V. Sharma, Lepton Photon 2011

- Arises in models with extra Higgs triplets
 - $\Phi^{++}, \Phi^+, \Phi^0$
- Triplet responsible for small neutrino mass
- Unknown neutrino mass matrix
 - unknown branching ratios → broad search
- Below $M \approx 2M_W$, only leptonic decays



CMS Preliminary

$$\text{BR}(\Phi^{++} \rightarrow e^+e^+) = 100\%$$

$$\text{BR}(\Phi^{++} \rightarrow e^+\mu^+) = 100\%$$

$$\text{BR}(\Phi^{++} \rightarrow \mu^+\mu^+) = 100\%$$

$$\text{BR}(\Phi^{++} \rightarrow e^+\tau^+) = 100\%$$

$$\text{BR}(\Phi^{++} \rightarrow \mu^+\tau^+) = 100\%$$

$$\text{BR}(\Phi^{++} \rightarrow \tau^+\tau^+) = 100\%$$

BP1: normal hierarchy


BP2: inverse hierarchy

BP3: degenerate masses



BP4: equal branchings

When $H^{++} \rightarrow l^+l^+$,
 $m_{H^{++}} > 250 - 300 \text{ GeV}$.


This bound cannot be applied
 when H^{++} does not decay into
 the same sign dilepton.



Branching ratio of H^{++}






$v_{\Delta} = 0.1 \text{ MeV}, m_{H^{++}} = 200 \text{ GeV}$



Chakrabarti, Choudhury, Godbole, Mukhopadhyaya, (1998);
 Chun, Lee, Park, (2003);
 Perez, Han, Huang, Li, Wang, (2008);
 Melfo, Nemevsek, Nesti, Senjanovic, (2011)

Phenomenology of $\Delta m \neq 0$ is drastically different from that of $\Delta m = 0$.

Branching ratios of H^+ , H and A



- ★ The $H^+ \rightarrow \phi^0 W^+$ mode can be dominant in the case of $\Delta m \neq 0$.
- ★ The $\phi^0 \rightarrow bb$ mode can be dominant when $v_\Delta > \text{MeV}$.

Constraints to extended Higgs sectors

- Non-minimal Higgs sectors

$SU(2)$ doublet + $\left[\begin{array}{l} \text{SU(2) singlets} \\ \text{SU(2) doublets} \\ \text{SU(2) triplets} \end{array} \right.$ Etc...

Ex. MSSM $\rightarrow$ 2HDM
Radiative seesaw models $\rightarrow$ 2HDM + sing
Type II seesaw model $\rightarrow$ triplet

There are many possibilities of non-minimal Higgs sectors.

- Constraints to extended Higgs sectors

- ★ Electroweak precision observables

- The rho parameter
- The W boson mass, ...

- ★ Flavor experiments


- Lepton flavor violation experiments ($\mu \rightarrow e\gamma$, $\mu \rightarrow eee$, ...)
- Quark flavor violation experiments ($b \rightarrow s\gamma$, K^0 - $\bar{K}^0$ mixing, ...)

In this talk, we focus on the constraint from the electroweak precision data.

Generation mechanisms for neutrino masses

Majorana masses of neutrinos are given by **the dimension 5 operator**, in which **2 units of lepton number are broken**.

$$\mathcal{L}_{\text{eff}} = \frac{c}{M} \bar{\nu}_L^c \nu_L \phi^0 \phi^0$$






The Higgs boson gets the VEV.

$$\frac{c \langle \phi^0 \rangle^2}{M} \bar{\nu}_L^c \nu_L$$


$$\langle \phi^0 \rangle \sim 246 \text{ GeV}$$

$$m_\nu \sim 0.1 \text{ eV}$$

$$c/M \sim 10^{-14} \text{ GeV}^{-1}$$



Dynamics of new physics models is mediated in the vertex.




What kind of NP models for generating neutrino masses are there ?

Seesaw Mechanism (Type-I)

Introducing right-handed neutrinos ν_R into the SM

$$\mathcal{L} = \mathcal{L}_{\text{SM}} - \underbrace{y_\nu \bar{L}_L \tilde{\Phi} \nu_R}_{\text{Dirac mass term}} - \frac{1}{2} \underbrace{m_R \bar{\nu}_R^c \nu_R}_{\text{Majorana mass term}} + \text{h.c.}$$


Dirac mass term **Majorana mass term**



$$m_\nu \sim \mathcal{O}(0.1) \text{ eV} \quad \Rightarrow \quad m_R \sim \mathcal{O}(10^{14}) \text{ GeV}$$


[In the case of $y_\nu = \mathcal{O}(1)$]

This model is simple but difficult to test.



$$\frac{c \langle \phi^0 \rangle^2}{M} \bar{\nu}_L^c \nu_L$$


$M \rightarrow m_R$



Extended Higgs sectors and neutrino masses

Tiny neutrino masses can be generated through dynamics of extended Higgs sectors at the TeV scale .


- ★ **Radiative seesaw models**: Neutrino masses can be generated at the loop level, where additional scalar bosons are running in the loop.
- ★ **Type-II seesaw model**: The Higgs triplet field is added to the SM.



These models are testable at the LHC

Radiative seesaw models

Neutrino masses are generated at the loop level.



Additional scalar bosons are running in the loop.

Neutrino mass

$$m_\nu = \left(\frac{1}{16\pi^2} \right)^N \frac{c' \langle \phi^0 \rangle^2}{M_\Phi}$$


Loop Factor

$$\frac{c \langle \phi^0 \rangle^2}{M} \bar{\nu}_L^c \nu_L$$

$$c \rightarrow \left(\frac{1}{16\pi^2} \right)^N c'$$

$$M \rightarrow M_\Phi$$

Loop factor $\times$
Yukawa couplings




Thanks for the loop factor, M_Φ (e.g., charged Higgs mass) can be taken to be TeV scale without finetuning in Yukawa couplings.

Variation for radiative seesaw models


Source of
L# violation

Scalar interaction


Zee Model (1980)




Zee-Babu Model (1986)




Ma Model (2006)



Krauss, Nasri, Trodden Model (2003)



Aoki, Kanemura, Seto Model (2009)



Majorana
mass

In the latter three models, a lightest Z_2 -odd particle can be a dark matter candidate.

Constraint to extended Higgs sectors

- There are two important experimental results.
 1. The electroweak ρ parameter is quite close to unity.
 2. FCNC processes are suppressed.

SU(2) doublet Higgs

- + Singlet [Both 1 and 2 are satisfied.]
- + Doublet [To satisfy 2, Z_2 symmetry is imposed]
- + Triplet [To satisfy both 1 and 2, parameter tuning is necessary.]

In this talk, we focus on the constraint from the electroweak precision measurement.